



REVIEW ARTICLE

Open Access

Growing burden of diabetes in Pakistan and the possible role of arsenic and pesticides

Haji Bahadar^{1,2}, Sara Mostafalou² and Mohammad Abdollahi^{1*}

Abstract

This review is undertaken to address the possible role of arsenic and pesticides in the prevalence of diabetes in Pakistan and to highlight a resourceful targeted research in this area.

A bibliographic search of scientific databases was conducted with key words of "epidemics of diabetes in Pakistan", "diabetes in Asia", "diabetes mellitus and environmental pollutants", "diabetes mellitus and heavy metals", "diabetes mellitus and pesticides", "prevalence of pesticides in Pakistan", and "heavy metals contamination of drinking water, "vegetables and fruits in Pakistan". More than 200 articles were examined. Studies reporting the prevalence of diabetes mellitus (DM), pesticides and heavy metal contamination of drinking water, fruits and vegetables were included in the study. According to WHO 2011 report, about 12.9 million people are suffering from DM and the number is constantly increasing. Water pollution is a major public health threat in Pakistan. Most of the people in Pakistan are exposed to arsenic and pesticides either in drinking water or through vegetables, fruits, and other edible items with various concentrations above the WHO/FAO permissible limits. Being an agricultural country, a 1169% increase has been recorded with the use of different types of pesticides since last two decades, and almost similar rise in the burden of diabetes.

There is a growing global concern of arsenic and pesticides exposure with the incidence of DM. Besides other factors, the environmental attributors in the incidence of DM in Pakistan have not been conclusively elucidated yet which in turn deserve a resourceful targeted research.

Keywords: Arsenic, Diabetes mellitus, Environmental pollutants, Heavy metals, Pakistan, Pesticides, Review

Introduction

Pollution is the addition of substances or energy to the environment, likely to produce harmful effects on human health and ecosystem. Heavy industrialization and scientific developments have led to the addition of detrimental chemicals to the environment in the form of heavy metals, agrochemicals, pesticides and hydrocarbons [1]. Environmental pollution has been a major health concern throughout the world since very long, but the risks are higher in underdeveloped countries. In low developed countries, environmental pollution contributes about 8-9% of total disease burden [2].

Pakistan, a country with a population exceeding 180 million, having four provinces along with federally

administered tribal areas, is the 6th populous country of the world with a very small economy [3].

Arsenic exposure in Pakistan

The presence of arsenic (As) in drinking water has become a major public health concern around the world. Arsenic has been recognized globally as the most toxic inorganic contaminant of drinking water [4]. Water sources of Asian countries, including Pakistan are among the most affected ones for As contamination [5,6].

Regional status of arsenic exposure in Pakistan

Sindh province

Approximately 36% people in Sindh province consume drinking water or vegetables containing As above the WHO limit [7]. The ground water sources in some rural areas in Sindh province contain As up to 1.1 mg/L [8]. A study published by Arain et al. reported the contamination of vegetables with As. The vegetables grown in the south

* Correspondence: Mohammad@TUMS.Ac.Ir

¹Department of Toxicology and Pharmacology, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, International Campus, Tehran University of Medical Sciences, Tehran, Iran

Full list of author information is available at the end of the article

east part of Sindh contain As in the range of 0.90-120 mg/kg, exceeding WHO/FAO limit of 0.0001 mg/kg [9]. Drinking water samples tested from districts Tharparkar, Matiari and Jamshoro have been shown to contain As in the range of 0.013-2.09 mg/L [7,10,11]. Data about the presence of As in either drinking water or vegetable from other areas of Sindh are not available to completely understand the status of peoples in term of As exposure in this province.

Punjab province

Punjab is the most populated province of Pakistan. The drinking water sources of Punjab province have been reported to contain As level above the WHO safe limit. According to reports, approximately 20% population in Punjab is exposed to As contaminated water [12]. Drinking water sources of some regions in Punjab, such as Bahawalpur, Gujranwala, Kasur, Lahore, Multan, Muzaffargarh, Rahim Yar Khan and Sheikhupura, have been reported to contain As level in the average range of 0.0794 to 0.9 mg/L [6,8,13-15]. "Pakistan Council of Research in Water Resources" a government body responsible for water quality in Pakistan, has reported As contents above the WHO permissible limits in drinking water sources of some other regions such as Attock, Multan, Sargodha and Bahawalpur. However, the values for As content in water samples of these areas have not been made public [16]. Overall, compared to other provinces of Pakistan, water samples of many areas of Punjab province have been studied for the presence of As contents. It has been reported that drinking water sources of Punjab province contain comparatively high As contents.

Khyber PakhtoonKhwa province

Arsenic from the soil gets its way to ground water and ultimately enters the crops. Beside presence of As in drinking water sources, it has been reported that wheat, which is the main edible crop of the Pakistani population, contain As contents above safe limits. Al-Othman et al. have studied wheat crop samples from different districts of Khyber PakhtoonKhwa for bioaccumulation of As contents. The results of this study showed that wheat crop in these areas contains As in the range of 0.005-1.113 mg/kg. The permissible limits of As in agronomic crops is 0.43 mg/kg [17]. However, WHO/FAO (Codex-1995 amended 2009) has not defined any permissible limit for As in the wheat crop Table 1.

Balochistan province

Studies about the presence of As prevalence in Balochistan province are limited. However, some studies have reported the presence of As within the range of permissible limits of WHO in some selected districts [13].

Table 1 Existence of As in drinking water sources of various regions of Pakistan

References	Sample studied	Arsenic	Result
[15]	Drinking water	As	79.4 µg /L
[8]	Drinking water	As	1.5-5 µg /L
[14]	Drinking water	As	Reported as "above WHO limit"
[6]	Drinking water	As	10-906 µg/L
[5]	Drinking water	As	906 µg/L
[11]	Drinking water	As	3-106 µg/L
[16]	Drinking water	As	Reported as "above WHO limit"
[18]	Drinking water	As	32-1900 µg/L
[17]	Wheat crop	As	0.005-1.113 mg/kg
[9]	Vegetables	As	0.90-120 mg/kg

Maximum residue limit for As in the vegetable is 0.0001 mg/kg (WHO/FAO).

Maximum residue limit of As for wheat crop is 0.43 mg/kg (Al-othman et al. (2012).

The permissible limit for As in drinking water is 10 µg/L (WHO 2006).

Pesticides exposure in Pakistan

Pesticides presence poses a great threat to the environment and human life. The use of pesticides in agriculture is substantially increasing from the last four decades for protection of crops [19,20]. It is astonishing to note that very small quantities of applied pesticides reaches the target organism accurately and major part of "applied pesticides" is dispersed in the environment and enters the human food chain [20]. Migration of applied pesticides in drinking water and subsequent entry to food chain has remained a global concern and several cases of drinking water contamination have been reported in the developed world [21]. Pakistan being an agriculture country consumes approximately 70 thousand tons of different pesticides annually and the use of different types of pesticides in Pakistan has increased by 1169% in the last 2 decades [22-24]. Each crop in Pakistan receives at least 10 different types of pesticides, which is an alarming signal for public health [23,25]. This huge application of pesticides in agriculture sector has led to the contamination of drinking water sources, vegetables, cattle food, milk, and fish samples throughout the country [26,27]. Some water sources of Punjab province have been reported by Tariq et al. to contain different types of pesticides like bifenthrin, cyhalothrin, carbofuran, endosulfan, methyl parathion and monocrotophos exceeding the permissible limits defined by WHO/FAO. Fruits and vegetable collected from various parts of the country and tested for the presences of different types of pesticides have been reported to contain pesticides like, carbofuran, dimethoate, deltamethrin, cypermethrin, and chlorpyrifos contents exceeding the WHO/FAO maximum residue limits [27-29]. Water sources of the cotton growing areas of, both Punjab and Sindh province, and tobacco growing areas of Khyber

PakhtoonKhwa have been found contaminated with pesticides. As Pakistan is an agricultural country, therefore a huge amount of different pesticide use for protection of crops is imminent. There are published few studies regarding the contamination of vegetables and other food items with pesticide residues from all parts of the country. Determination of pesticides in vegetables and drinking water, has remained an understudied subject in Pakistan. All available studies about the prevalence of pesticides in various areas have been accumulated in Table 2, Figure 1.

Evidences for association of As and pesticides with incidence of diabetes

Arsenic

Enough experimental and epidemiological evidences are available which suggest that As exposure add to the incidence of diabetes [33-36]. Navas-Acien et al. have published a cross sectional study of 788 individuals chronically exposed to As in the USA. The researchers found a strong correlation between chronic exposure to low level of As and non-insulin diabetes mellitus [37]. Exposure to As, either in drinking water or from

Table 2 Pesticides detected in water samples of various regions in Pakistan

Reference	Sample studied	Pesticides	(WHO/FAO) permissible limit	Result
[24]	Ground water	Bifenthrin	ND	11 µg/L
		Carbofuran	7 µg/L	36 µg/L
		Methyl parathion	9 µg/L	3 µg/L
		Monocrotophos	3 µg/L epa	20 µg/L
		Carbofuran	7 µg/L	36 µg/L
		Endosulfan	ND	6 µg/L
		Cyhalothrin	ND	7 µg/L
[30,31]	Ground water	Dichlorvos	ND	0.03-0.45 µg/L
		Mevinphos	ND	0.06-0.21 µg/L
		Dimethoate	6 µg/L	0.0-0.15 µg/L
		Methyl- parathion	ND	0.0-0.06 µg/L
		Chlorpyrifos	30	0.0-0.03 µg/L
		Fenitrothion	ND	0.0-0.2 µg/L
		Endosulfan	ND	0.0-0.2 µg/L
		Profenphos	3 µg/L	0.01-0.17 µg/L
		Carbofuran	7 µg/L	0.0-0.26 µg/L
		Lindane	2 µg/L	0.11 µg/L
[27]	Fruits	Cypermethrin	0.1 mg/kg	0.94 mg/kg
		Deltamethrin	ND	0.039 mg/kg
		Dimethoate	ND	0.139 mg/kg
		Endosulfan	ND	0.774 mg/kg
[32]	Vegetables	Lindane	0.5 mg/kg	4.21 mg/kg
		Cypermethrin	ND	1.63 mg/kg
	Luffa	Methylparathion	ND	1.71 mg/kg
		Methylparathion	0.2 mg/kg	2.5 mg/kg
		Methamidopos	1.0 mg/kg	2.60 mg/kg
	Cauliflower	P,P, DDT	1.0 mg/kg	10.3 mg/kg
		Methylparathion	ND	3.15 mg/kg
		Methamidopos	0.5 mg/kg	4.61 mg/kg
	Onion	Cypermethrin	0.1 mg/kg	1.8 mg/kg
		Malathion	3.0 mg/kg	10 mg/kg
Tomato	Fenvalerate	2.2 mg/kg	1.0 mg/kg	

ND: not defined.



Figure 1 Map representing areas with high contents of arsenic, pesticides and DM prevalence in Pakistan.




Figure 2 A Schematic illustration of possible pathways by which As and pesticides may induce diabetes.

other environmental sources, and its possible link with incidence of diabetes has been reported in epidemiologic studies published from various parts of the world like Bangladesh, Taiwan, South Korea, Cyprus, Serbia, China and Mexico [35,38-43]. Recently a population based study published from Iran has suggested a relationship between even a lower exposure level of As (20–30 µg/L) and incidence of diabetes and hypertension [44]. Recently a workshop was conducted by the U.S. National Toxicology Program on environmental chemicals and the incidence of DM in which various epidemiological and experimental evidences were evaluated. It has been concluded that studies regarding As exposure with an incidence of DM are suggestive but not sufficient and further objective research in this area is highly recommended [45].

Multiple mechanisms may be involved in As induced diabetes. In experimental studies As has been reported to act on multiple targets such as, affecting insulin sensitivity, altering β cells function, alteration of β cells signaling pathways, disturbing glucose production in the liver, and the reduction of insulin secretion and initiation of oxidative stress in the pancreas [46-49] Figure 2, Table 3. Evidences from population based and experimental studies mentioned above are suggestive enough to prove that chronic As exposure may lead to the incidence of diabetes.

Pesticides

Numerous experimental studies are available about the toxic effects of pesticides and the occurrence of type 2 diabetes. Organophosphorus and organochlorine types

Table 3 Evidences for association of As and pesticides with diabetes

Reference	Type of study	Compound	Doses	Duration	Outcome/result
[36]	Experimental	As	10 mg/kg	>3 months	Islets damage Insulin secretion ↓ reactive oxygen species (ROS) ↑
[50]	Experimental (Pancreatic β cells)	As	0.5-2 µM	>3 months	Insulin secretion ↓ Insulin mRNA levels ↓
[49]	Experimental	As	5.55 mg/kg	30 days	Liver glycogen level ↓ Blood sugar level ↑ Glutamate pyruvate transaminase activity ↓ Glucose 6-phosphatase activity ↓
[48]	Experimental	As	1 mM	–	PI-3 kinase independent; SAPK2/p38 IUF-1 translocation from cytoplasm to nucleus ↑
[35]	Epidemiologic	As	-	>3 months	Positive association of As exposure with diabetes
[37]	Epidemiologic	As	-	>3 months	Positive association of As exposure with diabetes
[38]	Epidemiologic	As	-	>3 months	Positive association of As exposure with diabetes
[51]	Epidemiologic	As	-	>3 months	Positive association of As exposure with diabetes
[39]	Epidemiologic	As	-	>3 months	Positive association of As exposure with diabetes
[41]	Epidemiologic	As	-	>3 months	Positive association of As exposure with diabetes
[42]	Epidemiologic	As	-	>3 months	Positive association of As exposure with diabetes
[52]	Epidemiologic	As	-	>3 months	Positive association of As exposure with diabetes
[43]	Epidemiologic	As	-	>3 months	Positive association of As exposure with diabetes
[44]	Epidemiologic	As	-	>3 months	Positive association of As with diabetes in population
[53]	Experimental	Malation	100-400 ppm	30 days	Hepatic glycogen phosphorylase ↑ Phosphoenol pyruvate carboxy kinase ↑
[54]	Experimental	Malation	25-100 mg/kg/day	32 days	Phosphoenol pyruvate carboxy kinase Glucose 6-phosphatase ↑
[55]	Epidemiologic	Organo -chlorine Pesticides	-	>3 months	Positive association of pesticides with diabetes Insulin resistance observed
[56]	Epidemiologic	Organo -chlorine Pesticides	-	>3 months	Positive association of pesticides with diabetes
[57]	Epidemiologic	Malation	-	>3 months	Positive association of pesticides with diabetes Insulin resistance observed




Figure 3 Trend in the rise of diabetic patients in Pakistan.

of pesticides in particular have been mentioned to possess deleterious effect on glucose metabolism and insulin secretion [53,55,57]. However, few epidemiologic studies exist in the literature about the role of pesticides in the occurrence of diabetes [58,59]. A prospective study published by Montgomery et al. have reported a high prevalence of diabetes in the 33457 licensed pesticide applicators in the US, and the ratio of diabetes incidence was noted more in organochlorine and organophosphorus type of pesticide applicators [56]. Pesticides may cause diabetes by affecting multiple pathways involved in glucose regulation. The involved mechanisms include oxidative stress, nitrosative stress, pancreatitis, inhibition of choline esterase, altered mitochondrial functions, and alteration of adrenal gland functions [54,60-62] (Figure 2, Table 3).

Overview of DM prevalence in Pakistan

DM is the leading chronic diseases and has emerged a big socioeconomic burden in Pakistan. Various population-based studies and national surveys have shown that DM is a highly prevalent disease in almost all regions of Pakistan, with an overall ratio of 22.04% in urban and 17.15% in rural areas [63-65]. According to various surveys, the pattern of DM prevalence in Pakistan is as Punjab; male, 16.6%, female, 19.3%, Khyber PakhtoonKhwa; 11.1% both sexes, Balochistan; 10.8% both sexes, Sindh; male 16.2% female 11.7% [66-68]. According to the latest data provided by the International Diabetes Federation, (www.idf.org).

(www.idf.org/Diabetesatlas/Data-Visualisations) Pakistan is among the leading countries with high prevalence of DM. As per WHO 2011 report total prevalence of diabetes is 12.9 million among them 9.4 million are diagnosed, while 38 million population is pre-diabetic and the number of diabetic patients would increase to 14 million by 2030 [69] Figures 3 and 4.

Discussion and conclusion

Like other poor countries, polluted water is also one of the major public health threats in Pakistan with special emphasis to drinking water. Pakistan stands at position 80 in the world community in term of drinking water quality. Water used for drinking purposes is contaminated with heavy metals and pesticides [22]. About 89% of drinking water sources have poor quality and are not according to the parameters set by WHO for human consumption [14,18]. Pakistan is an agricultural country, and for the sake of high crop yield, there has been recorded about 1169% increase in the use of different types of pesticides for the last two decades, and almost similar rise in the burden of diabetes, and according to various studies the number of diabetic patients in Pakistan would become two fold by 2030. According to World Bank Report 2010, Pakistan is facing health crises with high incidence of diabetes, affecting poor families which increase the burden of poverty (Figure 3).

Experimental and some epidemiologic studies gathered above suggest that environmental pollutants such as pesticides, As, Cd and Hg are risk factors in the incidence of DM. Excessive As presence has been reported in water sources of many other countries of the World like Argentina (1-9900 µg/L), China (0.05-850 µg/L), India (10-3200 µg/L), Brazil (0.5-350 µg/L), Mexico (8-620 µg/L), USA (1-100,000 µg/L), Taiwan (10-1820 µg/L), and particularly Bangladesh having 1-2500 µg/L [70-72]. And according published data As exposure has been considered as one of the risk factors in the incidence of DM in these countries and some countries like China, India, Brazil and USA are at the forefront of this epidemic (<http://www.idf.org/diabetesatlas/data-visualisations>).




Figure 4 Regional and gender wise prevalence of diabetes in Pakistan.

We are of the opinion that there might be a correlation of heavy metals like As and pesticide exposure with the prevalence of DM in Pakistan. Even though, there are many regions, where the drinking water sources are grossly contaminated with heavy metals and pesticides, but to correlate the prevalence of DM in exposed population and exposure to these pollutants; there is a complete lack of targeted epidemiologic research in the affected areas. We are of the view that along with other confounding factors, the role of As and pesticides in the epidemic of DM in Pakistan cannot be overlooked. Furthermore, there is a need of an urgent attention of concerned quarters like environmental researchers/toxicologists, governmental and non-governmental organizations to objectively evaluate the epidemics of DM in Pakistan from various dimensions including identifying the role of environmental pollutants and to take practical steps in this regard. No doubt, comprehensive and objective data derived from a resourceful targeted research on environmental pollutants and the prevalence of DM in Pakistan would be extremely laudable among environmental scientists. Data obtained from such targeted research will significantly add to the growing body of evidences about the role of environmental pollutants in the occurrence of chronic diseases.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

HB wrote the initial draft of the article and completed intensive literature review. SM edited the article. MA gave the idea, read, and edited the manuscript. All authors read and approved the final manuscript.

Funding sources

This paper is the outcome of a financially non supported in-house study.

Author details

¹Department of Toxicology and Pharmacology, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, International Campus, Tehran University of Medical Sciences, Tehran, Iran. ²Department of Pharmacology and Toxicology, School of Pharmacy, Ardebil University of Medical Sciences, Ardebil, Iran.

Received: 5 September 2014 Accepted: 27 November 2014

Published online: 14 December 2014

References

- Maduka H: Water pollution and man's health. *Internet J Gastroenterol* 2006, **4**(1):1-11.
- Briggs D: Environmental pollution and the global burden of disease. *Brit Med Bull* 2003, **68**(1):1-24.
- Waheed Y, Tahir S, Ahmad T, Qadri I: Sequence comparison and phylogenetic analysis of core gene of hepatitis C virus from Pakistani population. *Afr J Biotechnol* 2010, **9**(29):4561-4567.
- Chouhan S, Flora SJ: Arsenic and fluoride: two major ground water pollutants. *Indian J Exp Biol* 2010, **48**(7):666-678.
- Rahman MM, Naidu R, Bhattacharya P: Arsenic contamination in groundwater in the Southeast Asia region. *Environ Geochem Health* 2009, **31**(Suppl 1):9-21.
- Mukherjee A, Sengupta MK, Hossain MA, Ahamed S, Das B, Nayak B, Lodh D, Rahman MM, Chakraborti D: Arsenic contamination in groundwater: a global perspective with emphasis on the Asian scenario. *J Health Popul Nutr* 2006, **24**(2):142-163.
- Uqaili A, Mughal A, Maheshwari B: Arsenic contamination in ground water sources of district Matiari, Sindh. *Int J Chem Environ Eng* 2012, **3**(4):259-266.
- Haque IU, Nabi D, Baig M, Hayat W, Trefry M: Groundwater arsenic contamination: A multi-directional emerging threat to water scarce areas of Pakistan. *IAHS publication* 2008, **324**:24.
- Arain MB, Kazi TG, Baig JA, Jamali MK, Afridi HI, Shah AQ, Jalbani N, Sarfraz RA: Determination of arsenic levels in lake water, sediment, and foodstuff from selected area of Sindh, Pakistan: Estimation of daily dietary intake. *Food Chem Toxicol* 2009, **47**(1):242-248.
- Brahman KD, Kazi TG, Afridi HI, Naseem S, Arain SS, Ullah N: Evaluation of high levels of fluoride, arsenic species and other physicochemical parameters in underground water of two sub districts of Tharparkar, Pakistan: A multivariate study. *Water Res* 2013, **47**(3):1005-1020.
- Baig JA, Kazi TG, Arain MB, Afridi HI, Kandhro GA, Sarfraz RA, Jamal MK, Shah AQ: Evaluation of arsenic and other physico-chemical parameters of surface and ground water of Jamshoro. *Pakistan J Hazard Mater* 2009, **166**(2-3):662-669.
- Ahmad T, Kahlowan M, Tahir A, Rashid H: Arsenic an emerging issue: Experiences from Pakistan. Paper presented at: 30th WEDC international conference, Vientiane, Lao 2004. Water, Engineering and Developing Center, Loughborough University Leicestershire LE11 3TU UK: <http://wedc.lboro.ac.uk/resources/conference/30/Ahmad.pdf>.
- Soomro M, Khokhar M, Hussain W, Hussain M: *Drinking water Quality challenges in Pakistan*. Pakistan: Council of Research in Water Resources; 2011:17-28.
- PCRWR. National Water Quality Monitoring programme 5th Monitoring report 2005-2006: Pakistan Council for Research in Water Resources, Islamabad Pakistan. <http://www.pcrwr.gov.pk/publication%20pcrwr.aspx>.
- Ahmad SA, Gulzar A, Rehman HU, Soomro Z, Hussain M, Rehman M, Qadir M: Study of Arsenic in drinking water of district Kasur Pakistan. *World Appl Sci J* 2013, **24**(5):634-640.
- Tahir MA, Rasheed MH, Imran MS: Water Quality status in rural areas of Pakistan. *Pakistan Council of Research in Water Resources* 2010, (143):1-66. www.pcrwr.gov.pk/BtL_wr.../Rural%20Areas%20Report/WQSReport.pdf.
- Al-Othman ZA, Ali R, Al-Othman AM, Ali J A. Habila: Assessment of toxic metals in wheat crops grown on selected soils of Khyber Pukhtoon Khaw, Pakistan, irrigated by different water sources. *Arab J Chem* 2012. doi.org/10.1016/j.arabjc.2012.04.006
- Farooqi A, Masuda H, Siddiqui R, Naseem M: Sources of arsenic and fluoride in highly contaminated soils causing groundwater contamination in Punjab, Pakistan. *Arch Environ Contam Toxicol* 2009, **56**(4):693-706.
- Van der Werf HMG: Assessing the impact of pesticides on the environment. *Agric Ecosys Environ* 1996, **60**(2-3):81-96.
- Pimentel D: Amounts of pesticides reaching target pests: environmental impacts and ethics. *J Agric Environ Ethics* 1995, **8**(1):17-29.
- Caughy, M. E: Pesticides in surface waters: Distribution, trends, and governing factors. *Environmen Progress* 1999, (18(1):58-59 doi:10.1002/ep.670180107
- Azizullah A, Khattak MNK, Richter P, Häder DP: Water pollution in Pakistan and its impact on public health A review. *Environ Int* 2011, **37**(2):479-497.
- Tariq MI, Afzal S, Hussain I, Sultana N: Pesticides exposure in Pakistan: a review. *Environ Int* 2007, **33**(8):1107-1122.
- Tariq MI, Afzal S, Hussain I: Pesticides in shallow groundwater of Bahawalnagar, Muzafargarh, D.G. Khan and Rajan Pur districts of Punjab, Pakistan. *Environ Int* 2004, **30**(4):471-479.
- Khan MJ, Zia MS, Qasim M: Use of pesticides and their role in environmental pollution. *World Acad Sci Eng Technol* 2010, **72**:122-128.
- Hussain S, Masud T, Ahad K: Determination of pesticides residues in selected varieties of mango. *Pakistan J Nutr* 2002, **1**(1):41-42.
- Anwar T, Ahmad I, Tahir S: Determination of pesticide residues in fruits of Nawabshah district, Sindh, Pakistan. *Pak J Bot* 2011, **43**(2):1133-1139.
- Latif Y, Sherazi STH, Bhanger MI: Assessment of pesticide residues in commonly used vegetables in Hyderabad, Pakistan. *Ecotoxicol Environ Safety* 2011, **74**(8):2299-2303.
- Tahir S, Anwar T, Ahmad I, Aziz S, Mohammad A, Ahad K: Determination of pesticide residues in fruits and vegetables in Islamabad market. *J Environ Biol / Acad Environ Biol India* 2001, **22**(1):71-74.
- Ahad K, Anwar T, Ahmad I, Mohammad A, Tahir S, Aziz S, Baloch U: Determination of insecticide residues in groundwater of Mardan Division, NWFP, Pakistan: A case study. *Water SA* 2000, **26**(3):409-412.

31. Ahad K, Mohammad A, Khan H, Ahmad I, Hayat Y: **Monitoring results for organochlorine pesticides in soil and water from selected obsolete pesticide stores in Pakistan.** *Environ Monit Assess* 2010, **166**(1-4):191-199.
32. Masud SZ, Hasan N: **Pesticide Residues in Foodstuffs in Pakistan: Organochlorine, Organophosphorus and pyrethroid insecticides in Fruits and Vegetables.** In *Environ Toxicol Assess*. Edited by Richardson M. London: Taylor and Francis Ltd; 2004:270-280.
33. Huang CF, Chen YW, Yang CY, Tsai KS, Yang RS, Liu SH: **Arsenic and diabetes: current perspectives: Kaohsiung.** *J Med Sci* 2011, **27**(9):402-410.
34. Smith AH: **Arsenic and diabetes.** *Environ Health Perspect* 2013, **121**(3):A70-A71.
35. Rahman M, Tondel M, Ahmad SA, Axelsson O: **Diabetes mellitus associated with arsenic exposure in Bangladesh.** *Am J Epidemiol* 1998, **148**(2):198-203.
36. Yen CC, Lu FJ, Huang CF, Chen WK, Liu SH, Lin-Shiau SY: **The diabetogenic effects of the combination of humic acid and arsenic: in vitro and in vivo studies.** *Toxicol Lett* 2007, **172**(3):91-105.
37. Navas-Acien A, Silbergeld EK, Pastor-Barriuso R, Guallar E: **Arsenic exposure and prevalence of type 2 diabetes in US adults.** *JAMA J Am Med Asso* 2008, **300**(7):814-822.
38. Coronado-González JA, Del Razo LM, García-Vargas G, Sanmiguel-Salazar F, Escobedo-de la Peña J: **Inorganic arsenic exposure and type 2 diabetes mellitus in Mexico.** *Environ Res* 2007, **104**(3):383-389.
39. Wang SL, Chang FH, Liou SH, Wang HJ, Li WF, Hsieh DP: **Inorganic arsenic exposure and its relation to metabolic syndrome in an industrial area of Taiwan.** *Environ Int* 2007, **33**(6):805-811.
40. Pan WC, Seow WJ, Kile ML, Hoffman EB, Quamruzzaman Q, Rahman M, Mahiuddin G, Mostafa G, Lu Q, Christiani DC: **Association of low to moderate levels of arsenic exposure with risk of type 2 diabetes in Bangladesh.** *Am J Epidemiol* 2013, **178**(10):1563-1570.
41. Rhee SY, Hwang YC, Woo JT, Chin SO, Chon S, Kim YS: **Arsenic exposure and prevalence of diabetes mellitus in Korean adults.** *J Korean Med Sci* 2013, **28**(6):861-868.
42. Makris KC, Christophi CA, Pasi M, Ettinger AS: **A preliminary assessment of low level arsenic exposure and diabetes mellitus in Cyprus.** *BMC Public Health* 2012, **12**(1):334.
43. Wang W, Xie Z, Lin Y, Zhang D: **Association of inorganic arsenic exposure with type 2 diabetes mellitus: a meta-analysis.** *J Epidemiol Commu Health* 2014, **68**(2):176-184.
44. Mahram M, Shahsavari D, Oveisi S, Jalilolghadr S: **Comparison of hypertension and diabetes mellitus prevalence in areas with and without water arsenic contamination.** *J Res Med Sci* 2013, **18**(5):408-412.
45. Kuo C-C, Moon K, Thayer KA, Navas-Acien A: **Environmental chemicals and type 2 diabetes: an updated systematic review of the epidemiologic evidence.** *Curr Diab Rep* 2013, **13**(6):831-849.
46. Díaz-Villaseñor A, Burns AL, Hiriart M, Cebrián ME, Ostrosky-Wegman P: **Arsenic-induced alteration in the expression of genes related to type 2 diabetes mellitus.** *Toxicol Appl Pharmacol* 2007, **225**(2):123-133.
47. Tseng CH: **The potential biological mechanisms of arsenic-induced diabetes mellitus.** *Toxicol Appl Pharmacol* 2004, **197**(2):67-83.
48. Macfarlane WM, McKinnon CM, Felton-Edkins ZA, Cragg H, James RF, Docherty K: **Glucose stimulates translocation of the homeodomain transcription factor PDX1 from the cytoplasm to the nucleus in pancreatic beta-cells.** *J Biol Chem* 1999, **274**(2):1011-1016.
49. Pal S, Chatterjee AK: **Prospective protective role of melatonin against arsenic-induced metabolic toxicity in Wistar rats.** *Toxicology* 2005, **208**(1):25-33.
50. Díaz-Villaseñor A, Burns AL, Salazar AM, Sordo M, Hiriart M, Cebrián ME, Ostrosky-Wegman P: **Arsenite reduces insulin secretion in rat pancreatic β -cells by decreasing the calcium-dependent calpain-10 proteolysis of SNAP-25.** *Toxicol Appl Pharmacol* 2008, **231**(3):291-299.
51. Lai MS, Hsueh YM, Chen CJ, Shyu MP, Chen SY, Kuo TL, Wu MM, Tai TY: **Ingested inorganic arsenic and prevalence of diabetes mellitus.** *Am J Epidemiol* 1994, **139**(5):484-492.
52. Jovanovic D, Rasic-Milutinovic Z, Paunovic K, Jakovljevic B, Plavsic S, Milosevic J: **Low levels of arsenic in drinking water and type 2 diabetes in Middle Banat region, Serbia.** *Int J Hyg Environ Health* 2013, **216**(1):50-55.
53. Abdollahi M, Donyavi M, Pournourmohammadi S, Saadat M: **Hyperglycemia associated with increased hepatic glycogen phosphorylase and phosphoenolpyruvate carboxykinase in rats following subchronic exposure to malathion.** *Comp Biochem Physiol C Toxicol Pharmacol* 2004, **137**(4):343-347.
54. Mostafalou S, Eghbal MA, Nili-Ahmadabadi A, Baeri M, Abdollahi M: **Biochemical evidence on the potential role of organophosphates in hepatic glucose metabolism toward insulin resistance through inflammatory signaling and free radical pathways.** *Toxicol Indust Health* 2012, **28**(9):840-851.
55. Lee D-H, Lind PM, Jacobs DR, Salihovic S, van Bavel B, Lind L: **Polychlorinated Biphenyls and Organochlorine Pesticides in Plasma Predict Development of Type 2 Diabetes in the Elderly The Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS) study.** *Diab Care* 2011, **34**(8):1778-1784.
56. Montgomery M, Kamel F, Saldana T, Alavanja M, Sandler D: **Incident diabetes and pesticide exposure among licensed pesticide applicators: Agricultural health study, 1993-2003.** *Am J Epidemiol* 2008, **167**(10):1235-1246.
57. Raafat N, Abass MA, Salem HM: **Malathion exposure and insulin resistance among a group of farmers in Al-Sharkia governorate.** *Clin Biochem* 2012, **45**(18):1591-1595.
58. Saldana TM, Basso O, Hoppin JA, Baird DD, Knott C, Blair A, Alavanja MC: **Sandler D P: Pesticide exposure and self-reported gestational diabetes mellitus in the Agricultural Health Study.** *Diab Care* 2007, **30**(3):529-534.
59. Everett CJ, Matheson EM: **Biomarkers of pesticide exposure and diabetes in the 1999-2004 National Health and Nutrition Examination Survey.** *Environ Int* 2010, **36**(4):398-401.
60. Rahimi R, Abdollahi M: **A review on the mechanisms involved in hyperglycemia induced by organophosphorus pesticides.** *Pest Biochem Physiol* 2007, **88**(2):115-121.
61. Mostafalou S, Abdollahi M: **Pesticides and human chronic diseases: evidences, mechanisms, and perspectives.** *Toxicol Appl Pharmacol* 2013, **268**(2):157-177.
62. Saeidnia S, Abdollahi M: **Toxicological and pharmacological concerns on oxidative stress and related diseases.** *Toxicol Appl Pharmacol* 2013, **273**(3):442-455.
63. Shera AS, Jawad F, Maqsood A: **Prevalence of diabetes in Pakistan.** *Diab Res Clin Practice* 2007, **76**(2):219-222.
64. Akhter J: **The burden of diabetes in Pakistan: the national diabetes survey.** *J Pak Med Asso* 1999, **49**(9):205-205.
65. Zahid N, Claussen B, Hussain A: **Diabetes and impaired glucose tolerance in a rural area in Pakistan and associated risk factors.** *Diabetes Metab Syndr. Clin Res Reviews* 2008, **2**(2):125-130.
66. Shera A, Rafique G, Ahmed K, Baqai S, Khan I, King H: **Pakistan national diabetes survey prevalence of glucose intolerance and associated factors in North West Frontier Province (NWFP) of Pakistan.** *J Pak Med Asso* 1999, **49**(9):206-210.
67. Shera AS, Basit A, Fawwad A, Hakeem R, Ahmedani MY, Hydrie MZI, Khwaja IA: **Pakistan National Diabetes Survey: Prevalence of glucose intolerance and associated factors in the Punjab Province of Pakistan.** *Prim Care Diab* 2010, **4**(2):79-83.
68. Shera A, Rafique G, Khwaja I, Ara J, Baqai S, King H: **Pakistan national diabetes survey: prevalence of glucose intolerance and associated factors in Shikarpur, Sindh Province.** *Diab Med* 1995, **12**(12):1116-1121.
69. Shaw J, Sicree R, Zimmet P: **Global estimates of the prevalence of diabetes for 2010 and 2030.** *Diab Res Clin Practice* 2010, **87**(1):4-14.
70. Nordstrom DK: **Worldwide occurrences of arsenic in ground water.** *Science* 2002, **296**:2143-2145.
71. Mandal BK, Suzuki KT: **Arsenic round the world: a review.** *Talanta* 2002, **58**(1):201-235.
72. Chowdhury UK, Biswas BK, Chowdhury TR, Samanta G, Mandal BK, Basu GC, Chanda CR, Lodh D, Saha KC, Mukherjee SK: **Groundwater arsenic contamination in Bangladesh and West Bengal, India.** *Environ Health Perspect* 2000, **108**(5):393.

doi:10.1186/s40200-014-0117-y

Cite this article as: Bahadar et al.: Growing burden of diabetes in Pakistan and the possible role of arsenic and pesticides. *Journal of Diabetes & Metabolic Disorders* 2014 **13**:117.