

<sup>1</sup>Czech Technical University in Prague, Faculty of Biomedical Engineering, Prague, Czech Republic

<sup>2</sup>Charles University in Prague, 2nd Faculty of Medicine, Prague, Czech Republic

<sup>3</sup>University Hospital of North Norway, Norwegian Centre for Integrated Care and Telemedicine, Tromsø, Norway

<sup>4</sup>Charles University in Prague, 1st Faculty of Medicine, Prague, Czech Republic

### 389 THE IMPORTANCE OF PSYCHOLOGICAL ASSISTANCE AND SUPPORT FOR PEOPLE WITH DIABETES IN SOCIAL MEDIA

E. Patrakeeva<sup>1</sup>, T. Zagorovskaya<sup>2</sup>

<sup>1</sup>Saint-Petersburg Medical University, Endocrinology, Saint-Petersburg, Russia

<sup>2</sup>Saint Petersburg State University, Psychology, Saint-Petersburg, Russia

### 390 RELATIONSHIPS OF DIABETES BURNOUT (DB), DEPRESSION, ANXIETY, FREQUENCY OF SELF-MONITORING WITH HBA1C IN TYPE 1 DIABETES (T1D) PATIENTS—DATA FROM SOCIAL NETWORKS

E. Patrakeeva<sup>1</sup>, N. Novoselova<sup>2</sup>, L. Goncharova<sup>1</sup>, A. Zulevskaya<sup>1</sup>

<sup>1</sup>Saint-Petersburg Medical University, Endocrinology, Saint-Petersburg, Russia

<sup>2</sup>Federal Almazov North-West Medical Research Centre, Endocrinology, Saint-Petersburg, Russia

### 391 BENEFITS OF SOCIAL MEDIA POSTING INFORMATION FOR PATIENTS WITH DIABETES

E. Patrakeeva<sup>1</sup>, M. Dumicheva<sup>1</sup>

<sup>1</sup>Saint-Petersburg Medical University, Endocrinology, Saint-Petersburg, Russia

### 392 A DECISION SUPPORT TOOL FOR T2D MANAGEMENT EXPLOITING TEMPORAL DATA MINING ON HETEROGENEOUS DATA

L. Sacchi<sup>1</sup>, A. Dagliati<sup>1</sup>, V. Tibolla<sup>2</sup>, D. Segagni<sup>2</sup>, L. Chiovato<sup>3</sup>, R. Belluzzi<sup>4</sup>

<sup>1</sup>University of Pavia, Department of Electrical-Computer and Biomedical Engineering, Pavia, Italy

<sup>2</sup>IRCCS Fondazione Salvatore Maugeri, Laboratorio di Informatica e Sistemistica per la Ricerca Clinica, Pavia, Italy

<sup>3</sup>IRCCS Fondazione Salvatore Maugeri, Unità Operativa di Medicina Generale ed Endocrinologia, Pavia, Italy

### 393 EFFECT OF EDUCATIONAL INTERVENTION ON NUTRITIONAL BEHAVIOR AMONG PATIENTS WITH TYPE 2 DIABETES

D. Shojaei-zadeh<sup>1</sup>, N. Ghorbanzadeh<sup>2</sup>, M. Shojaei-zadeh<sup>2</sup>, N. Amini<sup>3</sup>, M. Shojaei-zadeh<sup>3</sup>

<sup>1</sup>School of Public Health- Tehran University of Medical Sciences, Department of Health Education & Promotion, Tehran, Iran

<sup>2</sup>Amirkabir University of Technology, Department of Biomedical, Tehran, Iran

<sup>3</sup>School of Public Health- Tehran University of Medical Sciences, Department of Public Health, Tehran, Iran

### 394 THE INFLUENCE OF HEALTH BELIEF MODEL COMPONENT ON SELF EFFICACY AMONG PATIENTS WITH TYPE 2 DIABETES

D. Shojaei-zadeh<sup>1</sup>, N. Ghorbanzadeh<sup>2</sup>, M. Shojaei-zadeh<sup>2</sup>, S. Vahidi<sup>3</sup>, M. Shojaei-zadeh<sup>1</sup>

<sup>1</sup>School of Public Health- Tehran University of Medical Sciences, Department of Health Education & Promotion, Tehran, Iran

<sup>2</sup>School of Public Health- Tehran University of Medical Sciences, Department of Public Health, Tehran, Iran

<sup>3</sup>Amirkabir University of Technology, Department of Biomedical, Tehran, Iran

<sup>4</sup>Shahid Beheshti University of Medical Sciences, Treatment Affairs, Tehran, Iran

### 395 WHATSAPP MESSENGER AS A TOOL TO IMPROVE GLUCOSE CONTROL IN TYPE 1 DIABETES PATIENTS ON PUMP THERAPY

F. Alkhalaf<sup>1</sup>, A. Elawaa<sup>1</sup>, A. Sabt<sup>1</sup>, F. Alyafei<sup>1</sup>, M. Elajamal<sup>1</sup>, Z. Mohammed<sup>1</sup>, H. Abubaker<sup>1</sup>, M. Itani<sup>1</sup>, N. El Sayed<sup>1</sup>, E. Mesaud<sup>1</sup>

<sup>1</sup>Hamad General Hospital, Pediatric Endocrinology/Diabetes, Doha, Qatar

### 396 EVOLUTION IN INSULIN NEEDS IN TYPE 2 DIABETIC (T2D) INSULIN-RESISTANT PATIENTS WITH AMBULATORY SUBCUTANEOUS INSULIN PUMP (AIP) - I RESIST' TRIAL - PRELIMINARY RESULTS

F. Traver<sup>1</sup>, S. Clave<sup>2</sup>, D. Huet<sup>3</sup>, E. Ghanassia<sup>4</sup>, S. Fendri<sup>5</sup>, V. Cosma<sup>6</sup>, P. Sérusclat<sup>7</sup>, J.P. Courrèges<sup>8</sup>

<sup>1</sup>Hôpital Bichat, Diabetology, Paris, France

<sup>2</sup>Centre Hospitalier Hôtel Dieu, Endocrinology and Diabetology, Le Creusot, France

<sup>3</sup>Hôpital Saint Joseph, Paris, France

<sup>4</sup>Clinique Sainte Thérèse, Endocrinology and Diabetology, Sète, France

<sup>5</sup>Centre Hospitalier Amiens, Amiens, France

<sup>6</sup>Centre Hospitalier Nîmes, Nîmes, France

<sup>7</sup>Clinique Les Portes du Sud, Vénissieux, France

<sup>8</sup>CHG Narbonne, Narbonne, France

### 397 THE "TECHNICAL DESK" OF THE DIABETES UNIT IN MILANO CA'GRANDA HOSPITAL: A VALUABLE HELP TO CLINICAL ACTIVITY

M. Bonomo<sup>1</sup>, B. Pintaudi<sup>1</sup>, O. Diotero<sup>2</sup>, G. Pizzi<sup>1</sup>, E. Mion<sup>1</sup>, F. Bertuzzi<sup>1</sup>

<sup>1</sup>A.O. Ospedale Niguarda Ca' Granda, Diabetes Unit, Milano, Italy

### 398 BOLUS WIZARD USE HAS A BENEFICIAL EFFECT IN GLYCEMIC CONTROL AMONG PEOPLE WITH DIABETES TYPE 1 ON INSULIN PUMP THERAPY

P. Giannoulaki<sup>1</sup>, I. Gountzioti<sup>2</sup>, F. Iludis<sup>3</sup>, A. Hatzioti<sup>3</sup>, A. Pappas<sup>4</sup>, D. Daskalaki<sup>4</sup>, T. Didangelos<sup>2</sup>

<sup>1</sup>AHEPA University General Hospital, Department of Nutrition and Dietetics, Thessaloniki, Greece

<sup>2</sup>Aristotle University of Thessaloniki- AHEPA University General Hospital, Diabetes Center- First Propedeutic Department of Internal Medicine, Thessaloniki, Greece