P11.05

Macracanthorhynchus ingens in long-eared hedgehog (*Hemiechinus auritus*): Histopathology and molecular characterization of the isolate

<u>Sanaz Jafarpoor Azami</u>¹, Eshrat Beigom Kia,¹, Iraj Mobedi¹, Zahra Heidari¹, Meysam Sharifdini², Kourosh Arzamani³, Bahareh Kamranrashani¹

¹Dept. of Medical Parasitology and Mycology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran, ²Department of Medical Microbiology, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran, ³Vector-borne Diseases Research Center, North Khorasan University of Medical Sciences, Bojnurd, Iran

Introduction: *Macracanthorhynchus ingens* belongs to perforated groups of acanthocephalan parasites which are usually parasite of raccoon, but they can also infect human and animals. This is the first report of *Macracanthorhynchus ingens* from long-eared hedgehog (*Hemiechinus auritus*) in Iran.

Methods: During examination of a hedgehog from Jajarm City in North Khorasan, Iran, nine acanthocephalan helminthes were found in small intestine. Helminthological clearing and staining techniques were applied and tissue sections prepared. For molecular analysis, after DNA extraction using designed primers for 18s ribosomal RNA, PCR amplification and sequencing were performed.

Results: Tissue sections revealed the penetration of proboscis in the intestine. The sequences showed 100% similarity with the sequences of *Macracanthorhynchus ingens* deposited in GenBank.

Conclusion: Occurrence of Macracanthorhynchus ingens in hedgehog in Iran constitutes new host species.